

Psychologisches Institut – Psychopathologie und Klinische Intervention (Universität Zürich)

On line terapije psihoterapijskog centra Sveučilišta u Zürichu se bave traumom i posttraumatskim stresnim poremećajem, teškim i produženim tugovanjem i poremećajima prilagodbe. Pristup je strukturiran i usmjeren na simptome i ima za cilj smanjiti smetnje i povećati cjelokupnu kvalitetu života. Terapija se sastoji od 12 do 16 seansi koje se provode dva puta tjedno. Osnova je kognitivna bihevioralna terapija čiji je cilj poticanje pacijenta da propituje svoje misli, prepoznaje disfunkcionalna ponašanja stvara bolje strategije za rješavanje teških životnih situacija.

Dodatne teme: Poremećaj prilagodbe, Trauma i posttraumatski stresni poremećaj, Produljeno žalovanje

Ciljana skupina	Tema	Vrsta dobre prakse	Zemlja
• Odrasli	• Depresija • Trauma	• Terapija	• Švicarska

Opis

The psychotherapeutic centre of the University of Zürich provides web-based therapies in the fields of trauma and post-traumatic stress disorders (PTSD), depression, persistent severe grief and adaptive disorders. The client is accompanied by a personal psychologist during the whole treatment. The therapy takes place via a secured internet platform and consists of 12 to 16 sessions. Each phase starts with a detailed introduction which explains the meaning of the practices. The client works on the tasks individually on the dates determined by him-or herself. The therapist reacts to the tasks within one work day. The basis of the modules is cognitive behavioural therapy which is scientifically proven to be appropriate for web-based therapies. The aim is to encourage the patient to question his/her thoughts, to recognize dysfunctional behaviours and to work out better strategies to cope with difficult life situations. The therapy for patients with trauma or post-traumatic stress disorder addresses people who had a traumatic experience at least three months ago and from which they still suffer. It consists of three phases which are self-confrontation, cognitive restructuring and sharing of fate. The therapy of depression includes five phases of treatment, these are: consideration of situations in which the patient is depressive, building up positive activities, become aware and change of negative and automatic thoughts, difficulties in interpersonal relationships as well as prevention of relapses. The treatment of persistent severe grief addresses people who sustained a traumatic loss and still suffer from that after six or more months. This kind of therapy is based on the same steps as the therapy of post-traumatic stress disorder. The treatments are not appropriate for people with addiction, dissociations, psychosis and suicidal tendencies.



FACHRICHTUNGEN

Home
Team
Studium
Forschung und Dissemination
Psychotherapie
Team
Trauma - Posttraumatische Belastungsstörungen (Traumatherapie)
Depression
Anhaltende schwere Trauer
Online-Therapie
Patienten-Information
Online-Therapie Behandlung
Alternativen
Wissenschaft
Anmeldung
Probleme im Alter
Leichte Alzheimer-Demenz
China-Projekt
Kontakt
Service public
Weiterbildung
Kontakt

Online-Therapie Behandlung

- Internet-Therapie von Trauma - Posttraumatischer Belastungsstörung
- Internet-Therapie von Depression
- Internet-Therapie von anhaltender schwerer Trauer

Die hier angebotene Internet-Behandlung von

- Trauma - Posttraumatische Belastungsstörung (PTBS)
- Depression
- Schwere Trauer

sind Therapien, die aus einer begrenzten Anzahl von 12 bis 16 Schreibsitzungen bestehen und auf vorgegebene Behandlungsschritte beruhen. Jeder Behandlungsphase geht eine ausführliche Einleitung voraus, die den Sinn einzelner Übungen erklärt. Die Aufgaben bearbeitet der Klient zu den selbst festgelegten Terminen zweimal wöchentlich. Sie werden vom Therapeuten innerhalb eines Arbeitstages beantwortet und kommentiert. Diese Interaktion findet innerhalb eines passwortgeschützten Internetportals statt, das strengsten Sicherheitsvorkehrungen genügt.

Es sind manualisierte, das heisst, strukturierte und auf Module aufbauende Therapien mit kognitiv-verhaltenstherapeutischer Grundlage, die sich in verschiedenen Studien für die Anwendung als Internet-Therapien als besonders geeignet erwiesen haben. Ziel der kognitiven Verhaltenstherapie ist es, ein besseres Verständnis und Einblick in die Erkrankung zu erhalten. Patienten werden ermutigt ihre problematischen Gedanken zu hinterfragen, hinderliche Verhaltensmuster zu erkennen und bessere Strategien zu erarbeiten um mit schwierigen Lebenssituationen umzugehen. Der Therapeut unterstützt den Klienten darin, korrektive, positive Lebenserfahrungen zu machen, wichtige Lebensziele wiederzufinden, sowie Massnahmen zu planen und durchzuführen.

Jezik originala

German

Zemlja

Switzerland

Poveznica

<http://www.psychologie.uzh.ch/de/fachrichtungen/psypath/Psychotherapie/Onlintherapie.html>

(access date: 07/04/2017)



CC - Attribution-NonCommercial-ShareAlike
<http://creativecommons.org/licenses/by-nc-sa/3.0/>

ILI. *Psychologisches Institut – Psychopathologie und Klinische Intervention (Universität Zürich)*. Skinuto sa Thu June 18, 2026 od <https://ecounselling4youth.gunet.gr/online-material/courses/TGP540/>



Co-funded by the
Erasmus+ Programme
of the European Union

The European Commission support for the production of this publication does not constitute an endorsement of the contents which reflects the views only of the authors, and the Commission cannot be held responsible for any use which may be made of the information contained therein

<https://www.ecounseling4youth.eu/project/>